## Seizure Emergency Care Plan and Medication Orders for School and Childcare Settings

PARENT/GUARDIAN complete a	nd sign tl	he top portion of form.		
Child Name:		Birth date:	1	
Parent/Guardian Contact:		Phone:	Place	
Emergency Contact:		Phone:	child's	
School:		Grade:	photo here	
Triggers: I tiredness I flashing lights I illness I hunger I temperature I Other:				
Seizure Aura (if any):				
Seizure history:				
Describe:				
Antiseizure Medication Taken at Home Common side effects				
Other Seizure Treatments/Special Diet Therapy:				
I give permission for school personnel to share this information, follow this plan, administer medication and care for my child and, if				
necessary, contact our health care provider. I assume full responsibility for providing the school with prescribed medication and devices. I approve this Seizure Emergency Care Plan for my child.				
	ing onnai			
			🗌 504 plan	
PARENT SIGNATURE D	ATE	SCHOOL NURSE SIGNATURE DATE		

## HEALTH CARE PROVIDER to complete all items, SIGN and DATE completed form.

IF YOU SEE THIS:	DO THIS:			
Convulsive Generalized Tonic Clonic: You will see loss of consciousness. Stiffening of the body. Rhythmic jerking movements. Convulsive seizures may last 1-5 minutes. The child may have a warning (aura) before the seizure. Sleepiness and confusion may occur after the seizure.	<ol> <li>Time the seizure</li> <li>Keep calm. Provide reassurance.</li> <li>Protect head, keep airway clear, turn on side if possible.</li> <li>Do not place anything in mouth.</li> <li>Call 911 if student is injured or has difficulty breathing.</li> <li>Call parent.</li> <li>Stay with student until recovered from seizure.</li> <li>Administer rescue treatments as marked below.</li> </ol>			
<b>Focal</b> : These seizures may begin with an aura. They may be partly alert or unconscious. You may see lip smacking, chewing, eye blinking, or picking at clothes. These seizures usually last 1-2 minutes.	<ol> <li>Time the seizure</li> <li>Gently guide child away from danger.</li> <li>Stay with student and reassure them until recovered from seizure.</li> <li>Do not treat staring that is stopped by a touch or a nudge.</li> <li>Call parent.</li> <li>Administer rescue treatments as marked below.</li> </ol>			
Absence: You will see quick changes in alertness. May see eye flutter or small twitching. Usually last less than 10 seconds.				
Rescue Treatments				
<ul> <li>Child has a VNS. Child/staff may swipe with aura. Staff may swipe at onset of seizure and every 60 seconds until seizure stops. Give rescue medications below if seizure does not stop withinminutes.</li> <li>If seizure lasts longer than minutes administer:</li> </ul>				
🗌 Diastatmg rectally 🛛 Midazola	ammg in the nose 🛛 🗌 Clonazepammg in the cheek			
Multistep seizure rescue plan – Please see attached letter for details.				
If <u>cluster</u> of or more seizures in min administer:				
Diastatmg rectally Didazolammg in the nose Clonazepammg in the cheek				
Multistep seizure rescue plan – Please see attached letter for details.				
If emergency medication is administered: 🔲 Call 911 immediately or 🗌 Call 911 if seizure does not stop within 5 minutes				
Other:				
If no emergency medication is at school and the child is experiencing seizures:				
Call family to bring medications to school or pick up child. Call EMS if seizure lasts more than min				
Accommodations: Always take seizure action plan and emergency medication for school activities, sports and field trips,				

**Accommodations**: Always take seizure action plan and emergency medication for school activities, sports and field trips. Close adult supervision when swimming or climbing.